

## Who can administer

Administration RESTRICTED - see [Appendix 1](#)

## Important information

- This drug may only be used under supervision of a **cardiologist in the Coronary Care unit**
- **Advanced cardio-pulmonary life-support systems must be immediately available** <sup>(ref 1)</sup>
- Ensure **isoprenaline injection** has been ordered and is ready for use if required- see separate monograph.
- Unlicensed medicine - not routinely stocked in GUH - reorder when used
- This monograph refers specifically to the use of the drug in the **diagnosis of Brugada syndrome**

## Available preparations

Gilurytmal injection 50mg per 10ml ampoule

## Reconstitution

Already in solution

**Draw up using a 5 micron filter needle**

## Infusion fluids

Not required-product ready for use

## Methods of intravenous administration

### **Slow intravenous injection (for the diagnosis of Brugada syndrome)**

- Administer each 10mg over one minute <sup>(ref 1)</sup> (see Dose below), followed by Sodium Chloride 0.9% flush<sup>Å</sup> <sup>(ref 3)</sup>

## Dose in adults

### **Ajmaline challenge in the diagnosis of Brugada syndrome** <sup>(ref 1,2)</sup>

- Administer 10mg every two minutes until termination indicated (see below) OR target dose of 1mg/kg (to a usual maximum total dose of 80mg (some references suggest maximum 100mg <sup>(ref 3)</sup>))
- Check ECG and observations after each bolus

### **Drug administration should be terminated if** <sup>(ref 1,2)</sup>

- Maximum dose reached, **OR**
- Diagnostic Brugada pattern in at least one ECG lead: J point or ST segment elevation of 2mm with coved ST elevation in more than one right precordial lead (positive result)
- Occurrence of ventricular ectopic beats or ventricular tachycardia
- Sinus arrest

- 2nd or 3rd degree AV nodal block
- Prolongation of QRS duration by more than 30%

## Monitoring

- Monitor for tissue damage if extravasated (low pH) <sup>(ref 3)</sup>
- Continue ECG monitoring for at least 1 hour afterwards or until ECG normalises or any adverse effects resolve <sup>(ref 3)</sup>
- During testing monitor blood pressure and pulse at 5 minute intervals until recovery is evident <sup>(ref 3)</sup>
- Consider also monitoring SaO<sub>2</sub> <sup>(ref 3)</sup>
- Serum sodium levels during administration should not exceed 145 to 150mmol/L <sup>(ref 3)</sup>

## Further information

- Contains propylene glycol <sup>(ref 3)</sup>

## Storage

- Store below 25°C

## References

SPC translated from German to English Dec 2010 (linked on Medusa IV website)

1: Ajmaline for the Diagnosis of Brugada Syndrome, Brighton and Sussex hospitals September 2023

2: Ajmaline test- Brugada syndrome. University Hospitals of Leicester 21/01/2022

3: Intravenous drug administration guide, Medusa , downloaded 18/03/2026

## Therapeutic classification

Class 1a antiarrhythmic